

Goldsmith, J.R. Epidemiological studies of radio-frequency radiation: current status and areas of concern. *The Science of the Total Environment* 180:3-8, 1996.

Gorbach, I.N. Changes in nervous system of individuals exposed to microradiowaves for long period of time. *Zdravookhraneniye Belorussii* 5:51-53, 1982. JPRS 81865, pp. 24-28.

Gordon, Z.V. The problem of the biological action of UHF. In *The Biological Action of Ultrahigh Frequencies*, A.A. Letavet and Z.V. Gordon, eds., Academy of Medical Sciences, Moscow, 1960, pp. 1-4.

Gordon, Z.V. Hygienic evaluation of the working conditions of workers with UHF generators. *Ibid.*, pp. 18-21.

Gordon, Z.V. and Lobanova, Y.A. Temperature reaction of animals under the influence of UHF. *Ibid.*, pp. 57-59.

Gordon, Z.V. Investigation of the blood pressure in rats (bloodless method) under the influence of UHF. *Ibid.*, pp. 64-67.

Gordon, Z.V., Roscin, A.V. and Byckov, M.S. Main directions and results of research conducted in the USSR on the biologic effects of microwaves. In *Biologic Effects and Health Hazards of Microwave Radiation: Proceedings of an International Symposium, Warsaw, 15-18 Oct., 1973*, P. Czerski et al., eds., pp. 22-35.

Gvozdkova, Z.M., Anan'ev, V.M., Zenina, I.N. and Zak, V.I. Sensitivity of the rabbit's central nervous system to a continuous superhigh-frequency electromagnetic field. *Bulletin of Experimental Biology and Medicine* 58(8):943-947.

67 Microwaving Our Planet

Gribbin, J. What's Wrong With Our Weather? the climatic threat of the 21st century. Scribner, New York, 1979.

Grin', A.N. Effects of microwaves on catecholamine metabolism in the brain. Vrachebnoye Delo 10:129-130, 1978. JPRS 72606, pp. 14-16.

Hallowell, C. Trouble in the lily pads. Time, Oct. 28, 1996, p. 87.

Hanson, M. What's the fuss about calcium? Interactions between calcium, free radicals and heavy metals and relevance to health. In Electromagnetic Hypersensitivity: Proceedings of the 2nd Copenhagen Conference, May 1995, J. Katajainen and B. Knave, eds., pp. 27-31.

Hanson, M. Do you feel rusty? Heavy Metal Bulletin 3(2)7-9+, 1996.

Hawk, K. Case Study in the Heartland. Butler, Pa., 1996.

Healer, J. Review of studies of people occupationally exposed to radio-frequency radiations. In Symposium Proceedings. Biological Effects and Health Implications of Microwave Radiation. Richmond, Va., S. Cleary, ed., Sept. 1969, pp. 90-97.

Hocking, B. and Gordon, I. Childhood leukemia risk increase associated with TV towers in North Sydney for exposures well below the present public safety standard. D.O.E. poster presentation, Palm Springs Workshop, 1995.

Huai, C. Assessment of health hazard and standard promulgation in China. Biological Effects and Dosimetry of Non-

ionizing Radiation, NATO Conference, Erice, Italy, 1981, pp. 627-644.

IEEE Standard for Safety Levels with Respect to Human Exposure to Radio Frequency Electromagnetic Fields, 3 kHz to 300 GHz. IEEE C95.1-1991.

Infante-Rivard, C. Electromagnetic field exposure during pregnancy and childhood leukaemia. *Lancet* 346:177, 1995.

Inglis, L.P. Why the double standard? A critical review of Russian work on the hazards of microwave radiation. 1970 IEEE International Symposium on Electromagnetic Compatibility, July 14-16, 1970, pp. 168-172.

Ismailov, E.S. Effects of microwaves on the nervous system. All-Union Conference on Problems of Biophysical Neurodynamics and General Biophysics, 1969:27-29. *JPRS* 75515, pp. 34-35.

Johansson, O. and Liu, P.-Y. "Electrosensitivity", "Electro-supersensitivity" and "screen dermatitis": preliminary observations from on-going studies in the human skin. In Proceedings of the Cost 244 Workshop on Electromagnetic Hypersensitivity, N. Leitgeb, ed., Oct. 1994, pp. 52-57.

Johnson, C.C. and Guy, A.W. Nonionizing electromagnetic wave effects in biological materials and systems. *Proceedings of the IEEE* 60(6):692-718, 1972.

Justeson, D.R. Behavioral and psychological effects of microwave radiation. *Bull. N.Y. Acad. Med.* 55(11): 1058-1078, 1979.

Kalnins, T., Krizbergs, R. and Romancuks, A. Measurement of the intensity of electromagnetic radiation from the Skrunda radio location station, Latvia. The Science of the Total Environment 180:51-56, 1996.

Kauppi, M. DNA injuries in electrically sensitive and CFS patients. Heavy Metal Bulletin 3(2):14, 1996.

Kauppi, M. The porphyrin link. Heavy Metal Bulletin 3(2):23+, 1996.

Kitsovskaya, I.A. Investigation of the interrelationships between the basic neural processes in rats under the influence of UHF of various intensities. In The Biological Action of Ultrahigh Frequencies, A.A. Letavet and Z.V. Gordon, eds., Academy of Medical Sciences, Moscow, 1960. JPRS 12471, pp. 75-82.

Klimkova-Deutschova, E. Neurologic findings in persons exposed to microwaves. In Biologic Effects and Health Hazards of Microwave Radiation: Proceedings of an International Symposium, Warsaw, 15-18 Oct., 1973, P. Czerski et al., eds., pp. 268-272.

Knave, B. Research reported in Forskning & Praktik, Apr. 1992, reprinted in Electrical Sensitivity News 1(5):4-5, 1996.

Knoppe, K.G. Parameters of UHF fields determining the hygienic evaluation of working conditions and the problems of their measurement. In The Biological Action of Ultrahigh Frequencies, A.A. Letavet and Z.V. Gordon, eds., Academy of Medical Sciences, Moscow, 1960. JPRS 12471, pp. 5-17.

- Ko, M., Sze, N. and Prather, M. Better protection of the ozone layer. *Nature* 367:505-508, 1994.
- Kolodynski, A.A. and Kolodynska, V.V. Motor and Psychological functions of school children living in the area of the Skrunda Radio Location Station in Latvia. *The Science of the Total Environment* 180:87-93, 1996.
- Kolomytkin, O., Yurinska, M., Zharikov, S., Kuznetsov, V. and Zharikova, A. Response of brain receptor systems to microwave energy exposure. In *On the Nature of Electromagnetic Field Interactions with Biological Systems*, A.H. Frey, ed., 1994, pp. 195-206.
- Kondra, P. A., Smith, W.K., Hodgson, G.C., Bragg, D.B., Gavora, J., Hamid, M.A.K. and Boulanger, R.J. Growth and reproduction of chickens subjected to microwave radiation. *Canadian Journal of Animal Science* 50:639-644, 1970.
- Kondra, P.A., Hamid, M.A. and Hodgson, G.C. Effects of microwave radiation on growth and reproduction of the stocks of chickens. *Canadian Journal of Animal Science* 52:317-320, 1972.
- Konradi, A. Effect of the orbital debris environment on the high-energy Van Allen proton belt. *Science* 242:1283-1286, 1988.
- Korbel Eakin, S. and Thompson, W.D. Behavioral effects of stimulation by UHF radio fields. *Psychological Reports* 17:595-602, 1965.

- Korbel, S.F. and Fine, H.L. Effects of low intensity UHF radio fields as a function of frequency. *Psychon. Sci.* 9(9):527-528, 1967.
- Kowalski, Z. and Indulski, J.A. The strategy of targetted health surveillance. II. Genetically determined susceptibility to chemical substances and other issues related to health surveillance. *Polish Journal of Occupational Medicine* 3(4):357-374, 1990.
- Krasny-Ergen, W. Point heating and mechanical effects of short waves. *Archives of Physical Therapy* 21:362-366, 1940.
- Krueger, W.F., Giarola, A.J., Bradley, J.W. and Shrekenhamer, A. Effects of electromagnetic fields on fecundity in the chicken. *Ann. N.Y. Acad. Sci.* 247:391-400, 1975.
- Kunjilwar, K.K. and Behari, J. Effect of amplitude-modulated RF radiation on cholinergic system of developing rats. *Brain Research* 601:321-324, 1993.
- Kupfer, A. The trouble with cellular. *Fortune*, Nov. 13, 1995, pp. 179-188.
- Lai, H. and Singh, N.P. Acute low-intensity microwave exposure increases DNA single-strand breaks in rat brain cells. *Bioelectromagnetics* 16:207-210, 1995.
- Leitgeb, N. Electromagnetic hypersensitivity. Quantitative assessment of an ill-defined problem. In *Proceedings of the Cost 244 Workshop on Electromagnetic Hypersensitivity*, N. Leitgeb, ed., Oct. 1994, pp. 68-74.

Lerner, E.J. Biological effects of electromagnetic fields.
IEEE Spectrum, May 1984, pp. 57-69.

Letavet, A.A. and Gordon, Z.V. The Biological Action of
Ultrahigh Frequencies. Academy of Medical Sciences,
Moscow, 1960. JPRS 12471.

Letavet, A.A. and Gordon, Z.V. Recommendations for conducting
preliminary and periodic medical examinations of workers
with UHF sources. Ibid., pp. 123-125.

Letavet, A.A. and Gordon, Z.V. Temporary sanitary regulations
in work with generators of centimeter waves. Ibid, pp. 126-
130.

Letavet, A.A. and Gordon, Z.V. Temporary instructions on
the method of measuring the power flux density of UHF
energy at the working positions. Ibid, pp. 131-133.

Levitt, B.B. Electromagnetic Fields: A Consumer's Guide to
the Issues and How to Protect Ourselves. Harcourt, Brace,
N.Y., 1995.

Liebesny, P. Athermic short wave therapy. Archives of
Physical Therapy, Dec. 1938, pp. 736-740.

Lin, J.C. Microwave Auditory Effects and Applications.
Charles C. Thomas, Springfield, 1978.

Lobanova, Y.A. and Gordon, Z.V. Investigation of the olfactory
sensitivity in persons subjected to the influence of UHF.
In The Biological Action of Ultrahigh Frequencies, A.A.
Letavet and Z.V. Gordon, eds., Academy of Medical Sciences,
Moscow, 1960. JPRS 12471, pp. 50-56.

73 Microwaving Our Planet

- Lobanova, Y.A. Survival and development of animals with various intensities and durations of the influence of UHF. Ibid, pp. 60-63.
- Lobanova, Y.A. and Tolgskaya, M.S. Change in the higher nervous activity and interneuron connections in the cerebral cortex of animals under the influence of UHF. Ibid, pp. 68-74.
- MacCracken, M.C., Budyko, M.I., Hecht, A.D. and Izrael, Y.A. Prospects for Future Climate: A Special US/USSR Report on Climate and Climate Change. 1990.
- Magone, I. The effect of electromagnetic radiation from the Skruna Radio Location Station on Spirodela polyrhiza (L.) Schleiden cultures. The Science of the Total Environment 180:75-80, 1996.
- Maietta, V. Iridium project beams into Tempe. Business Journal, July 26, 1996, pp. 1+.
- Maitland, G. and Thomas, J.R. Behavioral effects of daily and weekly 1 mW/cm^2 electromagnetic radiation (EMR) in rats. Bioelectromagnetics 1:203, 1980.
- Makhijani, A. and Gurney, K. Mending The Ozone Hole. Institute for Energy and Environmental Research, 1992.
- Mann, K. and Roschke, J. Effects of pulsed high-frequency electromagnetic fields on human sleep. Neuropsychobiology 33:41-47, 1996.
- Marha, K. Maximum admissible values of HF and UHF electromagnetic radiation at work places in Czechoslovakia. In Symposium Proceedings. Biological Effects and Health

- Implications of Microwave Radiation, Richmond, Va., Sept. 1969, S. Cleary, ed., pp. 188-191.
- Marha, K. Microwave radiation safety standards in Eastern Europe. IEEE Transactions on Microwave Theory and Techniques, Vol. MTT-19(2):165-168, 1971.
- Marino, A.A. Environmental electromagnetic energy and public health. In Modern Bioelectricity, A.A. Marino, ed., Dekker, N.Y., 1988.
- Markarov, G., Markarova, I., Zaslavsky, A. and Geles, U. Hypersensitivity to EMF, and the dependence of brain bio-electrical activity and general hemodynamics in cerebral asthenic (CA) patients, exposed to radioactive irradiation upon EMF 20-80 Hz effect. In Proceedings of the 2nd Copenhagen Conference on Electromagnetic Hypersensitivity, May 1995, J. Katajainen and B. Knave, eds., pp. 57-60.
- McRee, D.I. Review of Soviet/Eastern European research on health aspects of microwave radiation. Bull. N.Y. Acad. Med. 55(11):1133-1151, 1979.
- McRee, D.I. Soviet and Eastern European research on biological effects of microwave radiation. Proc. IEEE 68(1):84-91, 1980.
- Medici, R.G. Considerations for science: where has all the science gone? In Risk/Benefit Analysis: The Microwave Case, N.H. Steneck, ed., S.F. Press, 1982, pp. 177-196.
- Mickey, G.H. Electromagnetism and its effect on the organism. N.Y.S. Journal of Medicine, July 1, 1963, pp. 1935-1942.
- Microwave News, Nov./Dec. 1995, pp. 1+, report on cancer studies.
- Millar, H. Rockets for the rest of us. Wired, Sept. 1996, pp. 102-110.

Molina, M.J. and Rowland, F.S. Stratospheric sink for chlorofluoromethanes: chlorine atom-catalysed destruction of ozone. *Nature* 249:810-812, 1974

Morgan-Hughes, J.A., Darveniza, P., Kahn, S.N., Landon, D.N., Sherratt, R.M., Land, J.M. and Clark, J.B. A mitochondrial myopathy characterized by a deficiency in reducible cytochrome b. *Brain* 100:617-640, 1977.

Navakatikian, M.A. and Tomashevskaya, L.A. Phasic behavioral and endocrine effects of microwaves of nonthermal intensity. In *Biological Effects of Electric and Magnetic Fields*, D.O. Carpenter and S. Ayrapetyan, eds., Academic Press, N.Y. 1994, pp. 333-342.

National Council on Radiation Protection and Measurements. *Biological Effects and Exposure Criteria for Radio-frequency Electromagnetic Fields*. Report #86, Apr. 2, 1986.

Newell, R.E. Water vapour pollution in the stratosphere by the supersonic transporter? *Nature* 226:70-71, 1970.

New Scientist, 24 Aug. 1996. Are We Killing Astronomy? pp. 28-31.

Nikogosyan, S.V. Influence of UHF on the cholinesterase activity in the blood serum and organs in animals. In *The Biological Action of Ultrahigh Frequencies*, A.A. Letavet and Z.V. Gordon, eds., Academy of Medical Sciences, Moscow, 1960. JPRS 12471, pp. 83-88.

National Institute for Occupational Safety and Health. *Hazard Assessment of the Electronic Component Manufacturing Industry*. DHHS Publication #85-100, Washington, Feb. 1985.

Nuessle, V.D. and Holcomb, R.W. Will the SST pollute the stratosphere? *Science* 168:1562, 1970.

- Nutrition Reviews 46(4), 1988. Lactic acidosis and mitochondrial myopathy in a young woman, pp. 157-163.
- Ockerman, P. Study of electrosensitive persons reported in Goteborgs-Posten, June 7, 1996. Summarized in Heavy Metal Bulletin 3(2):14, and in L. Sodergren, 1996 EMF Diary.
- Olsen, R.G. and Hammer, W.C. Microwave-induced pressure waves in a model of muscle tissue. Bioelectromagnetics 1:45-54, 1980.
- Olsen, R.G. Evidence for microwave-induced acoustic resonances in biological material. Bioelectromagnetics 1:219, 1980.
- Orlova, A.A. The clinic of changes of the internal organs under the influence of UHF. In The Biological Action of Ultrahigh Frequencies, A.A. Letavet and Z.V. Gordon, eds., Academy of Medical Sciences, Moscow, 1960. JPRS 12471, pp. 30-35.
- Oscar, K.J. and Hawkins, T.D. Microwave alteration of the blood-brain barrier system of rats. Brain Research 126: 281-293, 1977.
- Ouellet-Hellstrom, R. and Stewart, W.F. Miscarriages among female physical therapists who report using radio- and microwave-frequency electromagnetic radiation. American Journal of Epidemiology 138:775-786, 1993.
- Park, C.G. and Helliwell, R.A. Magnetospheric effects of power line radiation. Science 200:727-730, 1978.
- Parshad, R. et al. Differential sensitivity of cultured human cells of two-tissue origin to killing by low-level

microwave radiation. In Biological Effects of Electropollution, S. Dutta and R. Millis, eds., Information Ventures, Phila., 1986, pp. 71-76.

Pazderova, J., Pickova, J. and Bryndova, V. Blood proteins in personnel of television and radio transmitting stations. In Biologic Effects and Health Hazards of Microwave Radiation: Proceedings of an International Symposium, Warsaw, 15-18 Oct., 1973, P. Czerski, ed., pp. 281-288.

Pervushin, V.Y. Changes occurring in the cardiac nervous apparatus due to the action of ultra-high-frequency field. Bull. Exper. Biol. Med. 43:734-740, 1957.

Prather, M.J., Garcia, M.M., Douglass, A.R., Jackman, C.H., Ko, M. and Sze, N.D. The Space Shuttle's impact on the stratosphere. Journal of Geophysical Research 95(D11): 18,583-18,590, 1990.

Presman, A.S. and Levitina, N.A. Nonthermal action of microwaves on cardiac rhythm, I. Bull. Exper. Biol. Med. 53(1):36-39, 1962.

Presman, A.S. and Levitina, N.A. Nonthermal action of microwaves on the rhythm of cardiac contractions in animals, II. Bull. Exper. Biol. Med. 53(2):154-157.

Presman, A.S. Electromagnetic Fields and Life. Plenum Press, N.Y., 1970.

Ray, S. and Behari, J. Physiologic changes in rats after exposure to low levels of microwaves. Radiation Research 123:199-202, 1990.

- Rea, W.J., Pan, Y., Fenyves, E.J., Sujisawa, I., Samadi, N. and Ross, G. Electromagnetic field sensitivity. *Journal of Bioelectricity* 10:241-256, 1991.
- Roberti, B., Heebels, G.H., Hendricx, J.C., de Greef, A.H. and Wolthius, O.L. Preliminary investigations of the effects of low-level microwave radiation on spontaneous motor activity in rats. *Ann. N.Y. Acad. Sci.* 247:417-424, 1975.
- Sadchikova, M.N. State of the nervous system under the influence of UHF. In *The Biological Action of Ultrahigh Frequencies*, A.A. Letavet and Z.V. Gordon, eds., Academy of Medical Sciences, Moscow, 1960, pp. 25-29.
- Sadchikova, M.N. Clinical manifestations of reactions to microwave irradiation in various occupational groups. In *Biologic Effects and Health Hazards of Microwave Radiation: Proceedings of an International Symposium, Warsaw, 15-18 Oct., 1973*, P. Czerski et al., eds., pp. 261-267.
- Sadchikova, M.N., Kharlamova, S.F., Shatskaya, N.N. and Kuznetsova, N.V. Significance of blood lipid and electrolyte disturbances in the development of some reactions to microwaves. *Gigiyena Truda i Professional'nyye Zabolevaniya* 2:38-39, 1980. JPRS 77393, pp. 37-39.
- Sagripanti, J. and Swicord, M.L. DNA structural changes caused by microwave radiation. *Int. J. of Rad. Biol.* 50(1):47-50, 1986.
- Salford, L.G., Brun, A., Persson, B.R. and Eberhardt, J. Experimental studies of brain tumour development during exposure to continuous and pulsed 915 MHz radiofrequency radiation. *Bioelectrochemistry and Bioenergetics* 30: 313-318, 1993.

Sarkar, S., Ali, S., Behari, J. Effect of low power microwave on the mouse genome: a direct DNA analysis. Mutation Research 320:141-147, 1994.

Savitz, D.A. and Calle, E.E. Leukemia and occupational exposure to electromagnetic fields: review of epidemiologic surveys. Journal of Occupational Medicine 29(1): 47-51, 1987.

Schmitz, P., Siegenthaler, J., Stager, C., Tarjan, D. and Bucher, J.B. Long-term exposure of young spruce and beech trees to 2450-MHz microwave radiation. The Science of the Total Environment 180:43-48, 1996.

Selga, T. and Selga, M. Response of *Pinus sylvestris* L. needles to electromagnetic fields. Cytological and ultrastructural aspects. The Science of the Total Environment 180:65-73, 1996.

Servantie, B., Servantie, A.M., Etienne, J. Synchronization of cortical neurons by a pulsed microwave field as evidenced by spectral analysis of EEG from the white rat. Ann. N.Y. Acad. Sci. 247:82-86, 1975.

Shandala, M.G. and Vinogradov, G.I. Immunological effects of microwave action. Gigiyena i Sanitariya 10:34-38, 1978. JPRS 72956, pp. 16-21.

Shandala, M.G., Dumanskii, U.D., Rudnev, M.I., Ershova, L.K. and Los, I.P. Study of nonionizing microwave radiation effects upon the central nervous system and behavior reactions. Environmental Health Perspectives 30:115-121, 1979.

Shandala, M.G., Rudnev, M.I., Stoyan, Y.F. and Vinogradov, G.I. Main directions of Soviet research on biological

effects of microwave radiation. Gigiyena i Sanitariya 10:4-7, 1981. JPRS 84221, pp. 75-80.

Shandala, M.G., Vinogradov, G.I., Rudnev, M.I. and Rudakova, S.F. Effects of chronic exposure to microwaves on certain indicators of cellular immunity. Radiobiologiya 23(4):544-546, 1983.

Sherry, S. High Tech and Toxics. Golden Empire Health and Planning Center, Sacramento, 1985.

Shutenko, O.I., Kozyarin, I.P. and Shvayko, I.I. Effects of superhigh frequency electromagnetic fields on animals of different ages. Gigiyena i Sanitariya 10:35-38, 1981. JPRS 84221, pp. 85-90.

Siekierzynski, M. A study of the health status of microwave workers. In Biologic Effects and Health Hazards of Microwave Radiation: Proceedings of an International Symposium, Warsaw, 15-18 Oct. 1973, P. Czerski et al., eds., pp. 273-280.

Sikorski, M. and Bielski, J. Disturbances of glucose tolerance in workers exposed to electromagnetic radiation. Medycyna Pracy 47(3):227-231, 1996.

Silverman, Charlotte. Epidemiologic approach to the study of microwave effects. Bull. N.Y. Acad. Med. 55(11):1166-1181, 1979.

Smirnova, M.I. and Sadchikova, M.N. Determination of the functional activity of the thyroid gland by means of radioactive iodine in workers with UHF generators. In The Biological Action of Ultrahigh Frequencies, A.A. Letavet and Z.V. Gordon, eds., Academy of Medical Sciences, Moscow, 1960. JPRS 12471, pp. 47-49.

Sodergren, L. 1996 EMF Diary. Goteborg, Sweden.

Sokolov, V.V. and Arievich, M.N. Changes in the blood under the influence of UHF on the organism. In The Biological Action of Ultrahigh Frequencies, A.A. Letavet and Z.V. Gordon, eds., Academy of Medical Sciences, Moscow, 1960, pp. 39-41

Solon, L.R. A local health agency approach to a permissible environmental level for microwave and radiofrequency radiation. Bull. N.Y. Acad. Med. 55(11):1251-1266.

Souder, W. An amphibian horror story. New York Newsday, Oct. 15, 1996, p. B19+.

Steneck, N.H., ed. Risk/Benefit Analysis: The Microwave Case. San Francisco Press, 1982.

Susskind, C. Testimony before the Committee on Commerce hearing on the Radiation Control for Health and Safety Act of 1967. Hearings Before the Committee on Commerce, United States Senate, 90th Congress, 2nd Session on S.2067, S.3211, and H.R. 10790 to provide for the protection of the public health from radiation emissions. Part 2. Serial No. 90-49, U.S. Government Printing Office, Washington, 1968, p. 720.

Szent-Gyorgyi, A. Molecules, electrons, and biology. Trans. N.Y. Acad. Sci. 31:334-340, 1969.

Szmigielski, S., Bielec, M., Lipski, S., and Sokolska, G. Immunologic and cancer-related aspects of exposure to low-level microwave and radiofrequency fields. In Modern Bioelectricity, A.A. Marino, ed., Dekker, N.Y., 1988, pp. 861-925.

Szmigielski, S. and Gil, J. Electromagnetic fields and neoplasms. In Electromagnetic Biointeraction, G. Franceschetti et al., eds., Plenum, N.Y., 1989, pp. 81-98.

Szmigielski, S. Cancer morbidity in subjects occupationally exposed to high frequency (radiofrequency and microwave) electromagnetic radiation. The Science of the Total Environment 180:9-17, 1996.

Szuba, M. and Szmigielski, S. Change in reaction time to auditory and visual signals differentiates individual responses to short-term exposure to ELF electric fields and direct current stimulation. In Proceedings of the Cost 244 Workshop on Electromagnetic Hypersensitivity, Graz, Austria, Oct. 1994, N. Leitgeb, ed., pp. 94-105.

Takashima, S., Onaral, B., and Schwan, H.P. Effects of modulated RF energy on the EEG of mammalian brains. Radiation and Environmental Biophysics 16:15-27, 1979.

Tanner, J.A. Effect of microwave radiation on birds. Nature 210:636, 1966.

Tanner, J.A., Romero-Sierra, C. and Davie, S.J. Non-thermal effects of microwave radiation on birds. Nature 216: 1139, 1967.

Tanner, J.A. and Romero-Sierra, C. Bird feathers as sensory detectors of microwave fields. In Biological Effects and Health Implications of Microwave Radiation, S. Cleary, ed., U.S. Dept. of HEW, Washington, pp. 185-187.

Tarricone, L., Cito, C. and D'Inzeo, G. Ach receptor channel's interaction with MW fields. Bioelectrochemistry and Bioenergetics 30:275-285, 1993.

83 Microwaving Our Planet

Teixeira-Pinto, A.A., Nejelski, L.L., Cutler, J.L. and Heller, J.H. The behavior of unicellular organisms in an electromagnetic field. *Experimental Cell Research* 20:548-564, 1960.

Tell, R.A. and Mantiply, E.D. Population exposure to VHF and UHF broadcast radiation in the United States. *Proc. IEEE* 68(1):4-12, 1980.

Thomas, Y., Litime, H., Belkadi, L., Beneviste, J., and Schiff, M. Electronic transmission of phorbol-myristate acetate to human neutrophils. Referenced in P. French, *The Interaction of Electromagnetic Fields with Biological Systems*, 1996.

Tofani, S., Agnesod, G., Ossola, P., Ferrini, S. and Bussi, R. Effects of continuous low-level exposure to radio-frequency radiation on intrauterine development in rats. *Health Physics* 51(4):489-499, 1986.

Tolgskaya, M.S., Gordon, Z.V. and Lobanova, Y.A. Morphological changes in experimental animals under the influence of pulse and continuous UHF. In *The Biological Action of Ultrahigh Frequencies*, A.A. Letavet and Z.V. Gordon, eds., Academy of Medical Sciences, Moscow, 1960. JPRS 12471, pp. 94-103.

Tolgskaya, M.S. and Gordon, Z.V. Changes in the receptor and interoreceptor apparatuses under the influence of UHF. *Ibid.*, pp. 104-109.

Trinos, M.S. Frequency of diseases of digestive organs in people working under conditions of combined effect of lead and SHF-range electromagnetic energy. *Gigiyena i Sanitariya* 9:93-94, 1982. JPRS 84221, pp. 23-26.

- Veyret, B., Bouthet, C., Deschaux, P., de Seze, R., Geffard, M., Joussot-Dubien, J., le Diraison, M., Moreau, J.-M. and Caristan, A. Antibody responses of mice exposed to low-power microwaves under combined pulse-and-amplitude modulation. *Bioelectromagnetics* 12:47-56, 1991.
- Wei, L. Y. A new theory of nerve conduction. *IEEE Spectrum*, Sept. 1966, pp. 123-127.
- Wieske, C.W. Human sensitivity to electric fields. In *Proceedings of the First National Biomedical Sciences Instrumentation Symposium*, Los Angeles, July 14-17, 1962. Reprinted in *Electrical Sensitivity News* 1(5): 1-4, 1996.
- Zaret, M.M. Selected cases of microwave cataract in man associated with concomitant annotated pathologies. In *Biologic Effects and Health Hazards of Microwave Radiation: Proceedings of an International Symposium*, Warsaw, 15-18 Oct., 1973, P. Czerski et al., eds., pp. 294-301.
- Zalyubovskaya, N.P. and Kiselev, R.I. Effect of radio waves of a millimeter frequency range on the body of man and animals. *Gigiyena i Sanitariya* 8:35-39, 1978. JPRS 72956, pp. 9-15.
- Zmyslony, M., Gadzicka, E., Szymczak, W. and Bortkiewicz, A. Evaluation of selected parameters of circulatory system function in various occupational groups exposed to high frequency electromagnetic fields. II. Electrocardiographic changes. *Medycyna Pracy* 47(3):241-252, 1996.

About the Author

Arthur Firstenberg is chairman of the Cellular Phone Taskforce, a citizens' group formed in response to the uncontrolled growth of the cellular phone industry. He was electrically injured in 1981 after three years in medical school at the University of California, Irvine. A holistic health practitioner, he is also an expert in the effects of technology upon the environment. He has been studying and writing about electromagnetic radiation for the past 15 years.

E86

Second World Congress on Electricity and Magnetism in Biology & Medicine, Bologna, Italy, June 8-13, 97.

similar to that caused by increased temperature, but without elevated temperature. Although details of the mechanism of interaction of EM fields with cells remain unknown, the induction of the stress response appears to be an appropriate cellular response to a stimulus that is not normally part of its environment, and this may well provide a clue to the mechanisms involved.

We thank the NCI, the US Department of Energy and the Heineman Foundation for their support.

MS-20-4

THE BIOLOGICAL EFFECTS OF MICROWAVE RADIATION. S. Kwee¹ and P. Raskmark². ¹Institute of Medical Biochemistry, University of Aarhus, DK-8000 Aarhus C, Denmark. ²Institute of Communication Technology, Aalborg University, DK-9220 Aalborg Ø, Denmark.

In recent times the use of mobile telephones has accelerated, resulting in an increasing exposure of the environment to weak radiofrequency (RF) fields, transmitted from these devices.

In previous work we showed that cell growth is affected by exposure to weak electromagnetic (ELF) fields¹. Consequently the next thing to investigate was, if EM fields generated by microwave radiation, would have a similar effect on cell proliferation.

The field was generated by signal simulation of the Global System for Mobile communications (GSM) of 960 MHz. Cell cultures, growing in microtiter plates, were exposed in a specially constructed chamber, a Transverse Electromagnetic (TEM) cell. The Specific Absorption Rate (SAR) values for each cell well were calculated for this exposure system².

Experiments were performed on cell cultures of transformed human epithelial amnion cells (AMA), which were exposed in the TEM cell to 960 MHz microwave fields at 3 different power levels. The cells, growing in microtiter plates in monolayer cultures, were exposed for 20, 30 or 40 min respectively. Cell proliferation was determined immediately before exposure and again after a 24 hours' growth period. It was found that cell growth in the exposed cells differed from that in the control and sham exposed cells and a decrease in cell growth was seen. Cell proliferation during the period following exposure not only varied with the various SAR levels, but also with the length of exposure time. On the other hand repeated periods of exposure did not seem to change the effects. There was a general linear correlation between power level and growth changes.

However, the exposure time required to obtain the maximum effect was not the same for the various power levels. It turned out that at low power level the maximum effect was first reached after a longer exposure time than at higher power level. A similar phenomenon was registered in our studies on ELF electromagnetic fields³. Here we found that there was a linear correlation between the length of exposure time to obtain maximum effect and the field strength. The explanation could be found in terms of "window" effects or adaptation.

SARs were 2.1, 0.21 & 0.021 mW/kg.

References:

1. S. Kwee and P. Raskmark: Changes in cell proliferation due to environmental non-ionizing radiation, 1. ELF electromagnetic fields. *Bioelectrochem. Bioenerg.* 36 (1995) 109.
2. K. V. Steffensen, P. Raskmark and G. F. Pedersen: FDTD calculations of the EM-field distribution in a microtiter suspension well. *Proceedings of the 244 COST meeting on "Biological effects relevant to amplitude modulated RF fields"*. Kuopio, 1995.
3. S. Kwee and P. Raskmark: The minimizing effect of electromagnetic noise on the changes in cell proliferation caused by ELF electromagnetic fields. *Proceedings of the 3rd International Congress of the European Bioelectromagnetics Association*. Nancy 1996.

MS-20-5

USE OF NON-MAMMALIAN MODEL SYSTEMS TO INVESTIGATE MECHANISMS MEDIATING BIOLOGICAL INTERACTIONS WITH ELECTROMAGNETIC FIELDS. D.M. Binnering. Department of Biological Sciences, Florida Atlantic University, Boca Raton, Florida 33431, USA.

The question of whether power frequency (60 Hz) electromagnetic fields (EMF) present a health risk to humans remains highly controversial. While a variety of biological responses to EMF under laboratory conditions have been reported, the molecular mechanisms that mediate these phenomena remain elusive. Insight into the biochemical and genetic mechanisms that transduce EMF into biologically responsive signals is critical for clarifying the putative role of EMF in the development of certain human cancers.

The yeast *Saccharomyces cerevisiae* is an ideal model organism for studying molecular mechanisms of eukaryotic gene expression. This single-celled fungus is highly amenable to traditional genetic techniques. Additionally, recombinant DNA techniques are available for yeast which are not yet technically feasible with more complex cells, especially those of humans. While completion of the Human Genome Project remains at least 5-10 years away, the DNA sequence of the entire yeast genome is now available. Thus, experimental approaches can be designed using yeast which are not available for the other model organisms typically used in EMF studies.

Has studying yeast actually contributed to our understanding of cancer in humans? An unexpected outcome of cloning human genes has been the discovery that many cancer-related genes are very similar in all eukaryotes—from yeast to humans. Numerous human genes involved in regulating cell growth and are defective in human cancers were first identified as CDC (cell division control) genes in yeast. First identified in genetic studies using yeast, molecular clones of the yeast genes were then used to identify the homologous genes in humans. There is an ever-increasing list of human cancer genes that have structural and functional homologies in yeast.

Reported by Dr. Henry
who attended the workshop
of UNW. of WASH., Seattle, WA

The characteristic and discrete energies $h\nu$ found as emission and absorption of electromagnetic radiation by atoms and molecules extend to X-ray energies. As high-energy electrons strike the piece of metal in an X-ray tube, electrons are knocked out of the inner energy shell of the atoms. These vacancies are then filled by electrons from the second or third shell; emitted in the process are X rays having $h\nu$ values that correspond to the energy differences of the shells. One therefore observes not only the continuous spectrum of the bremsstrahlung discussed above but also X-ray emissions of discrete energies $h\nu$ that are characteristic of the specific elemental composition of the metal struck by the energetic electrons in the X-ray tube.

Trans-
parency of
water to
visible light

The discrete electromagnetic radiation energies $h\nu$ emitted or absorbed by all substances reflect the discreteness of the internal energies of all material things. This means that window glass and water are transparent to visible light; they cannot absorb these visible light quanta because their internal energies are such that no energy difference between a higher and a lower internal state matches the energy $h\nu$ of visible light. Figure 3 shows as an example the coefficient of absorption of water as a function of frequency ν of electromagnetic radiation. Above the scale of frequencies, the corresponding scales of photon energy $h\nu$ and wavelength λ are given. An absorption coefficient $\alpha = 10^{-4} \text{ cm}^{-1}$ means that the intensity of electromagnetic radiation is only one-third its original value after passing through 100 metres of water. When $\alpha = 1 \text{ cm}^{-1}$, only a layer one centimetre thick is needed to decrease the intensity to one-third its original value, and, for $\alpha = 10^3 \text{ cm}^{-1}$, a layer of water having a thickness of this page is sufficient to attenuate electromagnetic radiation by that much. The transparency of water to visible light, marked by the vertical dashed lines, is a remarkable feature that is significant for life on Earth.




Figure 3: The absorption coefficient for liquid water as a function of frequency. Also shown as abscissas are an energy scale (arrows) and a wavelength scale (vertical lines). The visible region of the frequency spectrum is indicated by the vertical dashed lines. The absorption coefficient for seawater is denoted by the dashed diagonal line at the left. The scales are logarithmic in both directions.

All things look so different and have different colours because of their different sets of internal discrete energies, which determine their interaction with electromagnetic radiation. The words *looking* and *colours* are associated with the human detectors of electromagnetic radiation, the eyes. Since there are instruments available for detecting electromagnetic radiation of any frequency, one can imagine that things "look" different at all energies of the spectrum because different materials have their own characteristic sets of discrete internal energies. Even the nuclei of atoms are composites of other elementary particles

and thus can be excited to many discrete internal energy states. Since nuclear energies are much larger than atomic energies, the energy differences between internal energy states are substantially larger, and the corresponding electromagnetic radiation quanta $h\nu$ emitted or absorbed when nuclei change their energies are even bigger than those of X rays. Such quanta given off or absorbed by atomic nuclei are called gamma rays (see *The electromagnetic spectrum* above).

PROPERTIES AND BEHAVIOUR

Scattering, reflection, and refraction. If a charged particle interacts with an electromagnetic wave, it experiences a force proportional to the strength of the electric field and thus is forced to change its motion in accordance with the frequency of the electric field wave. In doing so, it becomes a source of electromagnetic radiation of the same frequency, as described in the previous section. The energy for the work done in accelerating the charged particle and emitting this secondary radiation comes from and is lost by the primary wave. This process is called scattering.

Since the energy density of the electromagnetic radiation is proportional to the square of the electric field strength and the field strength is caused by acceleration of a charge, the energy radiated by such a charge oscillator increases with the square of the acceleration. On the other hand, the acceleration of an oscillator depends on the frequency of the back-and-forth oscillation. The acceleration increases with the square of the frequency. This leads to the important result that the electromagnetic energy radiated by an oscillator increases very rapidly—namely, with the square of the square or, as one says, with the fourth power of the frequency. Doubling the frequency thus produces an increase in radiated energy by a factor of 16.

This rapid increase in scattering with the frequency of electromagnetic radiation can be seen on any sunny day: it is the reason the sky is blue and the setting Sun is red. The higher-frequency blue light from the Sun is scattered much more by the atoms and molecules of the Earth's atmosphere than is the lower-frequency red light. Hence the light of the setting Sun, which passes through a thick layer of atmosphere, has much more red than yellow or blue light, while light scattered from the sky contains much more blue than yellow or red light.

The process of scattering, or reradiating part of the electromagnetic wave by a charge oscillator, is fundamental to understanding the interaction of electromagnetic radiation with solids, liquids, or any matter that contains a very large number of charges and thus an enormous number of charge oscillators. This also explains why a substance that has charge oscillators of certain frequencies absorbs and emits radiation of those frequencies.

When electromagnetic radiation falls on a large collection of individual small charge oscillators, as in a piece of glass or metal or a brick wall, all of these oscillators perform oscillations in unison, following the beat of the electric wave. As a result, all the oscillators emit secondary radiation in unison (or coherently), and the total secondary radiation coming from the solid consists of the sum of all these secondary coherent electromagnetic waves. This sum total yields radiation that is reflected from the surface of the solid and radiation that goes into the solid at a certain angle with respect to the normal (i.e., a line perpendicular to) the surface. The latter is the refracted radiation that may be attenuated (absorbed) on its way through the solid.

Superposition and interference. When two electromagnetic waves of the same frequency superpose in space, the resultant electric and magnetic field strength of any point of space and time is the sum of the respective fields of the two waves. When one forms the sum, both the magnitude and the direction of the fields need be considered, which means that they sum like vectors. In the special case when two equally strong waves have their fields in the same direction in space and time (i.e., when they are in phase), the resultant field is twice that of each individual wave. The resultant intensity, being proportional to the square of the field strength, is therefore not two but four times the intensity of each of the two superposing waves.

Signifi-
cance
of the
scattering
process

E86

Second World Congress for Electricity and Magnetism in Biology & Medicine, Bologna, Italy. June 8-13, 97.

similar to that caused by increased temperature, but without elevated temperature. Although details of the mechanism of interaction of EM fields with cells remain unknown, the induction of the stress response appears to be an appropriate cellular response to a stimulus that is not normally part of its environment, and this may well provide a clue to the mechanisms involved.

We thank the NCI, the US Department of Energy and the Heineman Foundation for their support.

MS-20-4

THE BIOLOGICAL EFFECTS OF MICROWAVE RADIATION. S. Kwee¹ and P. Raskmark². ¹Institute of Medical Biochemistry, University of Aarhus, DK-8000 Aarhus C, Denmark. ²Institute of Communication Technology, Aalborg University, DK-9220 Aalborg Ø, Denmark.

In recent times the use of mobile telephones has accelerated, resulting in an increasing exposure of the environment to weak radiofrequency (RF) fields, transmitted from these devices.

In previous work we showed that cell growth is affected by exposure to weak electromagnetic (ELF) fields¹. Consequently the next thing to investigate was, if EM fields generated by microwave radiation, would have a similar effect on cell proliferation.

The field was generated by signal simulation of the Global System for Mobile communications (GSM) of 960 MHz. Cell cultures, growing in microtiter plates, were exposed in a specially constructed chamber, a Transverse Electromagnetic (TEM) cell. The Specific Absorption Rate (SAR) values for each cell well were calculated for this exposure system².

Experiments were performed on cell cultures of transformed human epithelial amnion cells (AMA), which were exposed in the TEM cell to 960 MHz microwave fields at 3 different power levels. The cells, growing in microtiter plates in monolayer cultures, were exposed for 20, 30 or 40 min respectively. Cell proliferation was determined immediately before exposure and again after a 24 hours' growth period. It was found that cell growth in the exposed cells differed from that in the control and sham exposed cells and a decrease in cell growth was seen. Cell proliferation during the period following exposure not only varied with the various SAR levels, but also with the length of exposure time. On the other hand repeated periods of exposure did not seem to change the effects. There was a general linear correlation between power level and growth changes.

However, the exposure time required to obtain the maximum effect was not the same for the various power levels. It turned out that at low power level the maximum effect was first reached after a longer exposure time than at higher power level. A similar phenomenon was registered in our studies on ELF electromagnetic fields³. Here we found that there was a linear correlation between the length of exposure time to obtain maximum effect and the field strength. The explanation could be found in terms of "window" effects or adaptation.

SARs were 2.1, 0.21 & 0.021 mW/kg.

References:

1. S. Kwee and P. Raskmark: Changes in cell proliferation due to environmental non-ionizing radiation, 1. ELF electromagnetic fields. *Bioelectrochem. Bioenerg.* 36 (1995) 109.
2. K. V. Steffensen, P. Raskmark and G. F. Pedersen: FDTD calculations of the EM-field distribution in a microtiter suspension well. *Proceedings of the 244 COST meeting on "Biological effects relevant to amplitude modulated RF fields"*. Kuopio, 1995.
3. S. Kwee and P. Raskmark: The minimizing effect of electromagnetic noise on the changes in cell proliferation caused by ELF electromagnetic fields. *Proceedings of the 3rd International Congress of the European Bioelectromagnetics Association*. Nancy 1996.

MS-20-5

USE OF NON-MAMMALIAN MODEL SYSTEMS TO INVESTIGATE MECHANISMS MEDIATING BIOLOGICAL INTERACTIONS WITH ELECTROMAGNETIC FIELDS. D.M. Binninger. Department of Biological Sciences, Florida Atlantic University, Boca Raton, Florida 33431, USA.

The question of whether power frequency (60 Hz) electromagnetic fields (EMF) present a health risk to humans remains highly controversial. While a variety of biological responses to EMF under laboratory conditions have been reported, the molecular mechanisms that mediate these phenomena remain elusive. Insight into the biochemical and genetic mechanisms that transduce EMF into biologically responsive signals is critical for clarifying the putative role of EMF in the development of certain human cancers.

The yeast *Saccharomyces cerevisiae* is an ideal model organism for studying molecular mechanisms of eukaryotic gene expression. This single-celled fungus is highly amenable to traditional genetic techniques. Additionally, recombinant DNA techniques are available for yeast which are not yet technically feasible with more complex cells, especially those of humans. While completion of the Human Genome Project remains at least 5-10 years away, the DNA sequence of the entire yeast genome is now available. Thus, experimental approaches can be designed using yeast which are not available for the other model organisms typically used in EMF studies.

Has studying yeast actually contributed to our understanding of cancer in humans? An unexpected outcome of cloning human genes has been the discovery that many cancer-related genes are very similar in all eukaryotes—from yeast to humans. Numerous human genes involved in regulating cell growth and are defective in human cancers were first identified as CDC (cell division control) genes in yeast. First identified in genetic studies using yeast, molecular clones of the yeast genes were then used to identify the homologous genes in humans. There is an ever-increasing list of human cancer genes that have structural and functional homologies in yeast.

Reported by Dr. Henry ^{Hai} who attended the workshop * of Univ. of WASH., Seattle, WA

E86

Second World Congress on Electricity and Magnetism in Biology & Medicine, Bologna, Italy. June 8-13, 97.

similar to that caused by increased temperature, but without elevated temperature. Although details of the mechanism of interaction of EM fields with cells remain unknown, the induction of the stress response appears to be an appropriate cellular response to a stimulus that is not normally part of its environment, and this may well provide a clue to the mechanisms involved.

We thank the NCI, the US Department of Energy and the Heineman Foundation for their support.

MS-20-4

THE BIOLOGICAL EFFECTS OF MICROWAVE RADIATION. S. Kwee¹ and P. Raskmark². ¹Institute of Medical Biochemistry, University of Aarhus, DK-8000 Aarhus C, Denmark. ²Institute of Communication Technology, Aalborg University, DK-9220 Aalborg Ø, Denmark.

In recent times the use of mobile telephones has accelerated, resulting in an increasing exposure of the environment to weak radiofrequency (RF) fields, transmitted from these devices.

In previous work we showed that cell growth is affected by exposure to weak electromagnetic (ELF) fields¹. Consequently the next thing to investigate was, if EM fields generated by microwave radiation, would have a similar effect on cell proliferation.

The field was generated by signal simulation of the Global System for Mobile communications (GSM) of 960 MHz. Cell cultures, growing in microtiter plates, were exposed in a specially constructed chamber, a Transverse Electromagnetic (TEM) cell. The Specific Absorption Rate (SAR) values for each cell well were calculated for this exposure system².

Experiments were performed on cell cultures of transformed human epithelial amnion cells (AMA), which were exposed in the TEM cell to 960 MHz microwave fields at 3 different power levels. The cells, growing in microtiter plates in monolayer cultures, were exposed for 20, 30 or 40 min respectively. Cell proliferation was determined immediately before exposure and again after a 24 hours' growth period. It was found that cell growth in the exposed cells differed from that in the control and sham exposed cells and a decrease in cell growth was seen. Cell proliferation during the period following exposure not only varied with the various SAR levels, but also with the length of exposure time. On the other hand repeated periods of exposure did not seem to change the effects. There was a general linear correlation between power level and growth changes.

However, the exposure time required to obtain the maximum effect was not the same for the various power levels. It turned out that at low power level the maximum effect was first reached after a longer exposure time than at higher power level. A similar phenomenon was registered in our studies on ELF electromagnetic fields³. Here we found that there was a linear correlation between the length of exposure time to obtain maximum effect and the field strength. The explanation could be found in terms of "window" effects or adaptation.

SARs were 2.1, 0.21 & 0.021 mW/kg.

References:

1. S. Kwee and P. Raskmark: Changes in cell proliferation due to environmental non-ionizing radiation, 1. ELF electromagnetic fields. *Bioelectrochem. Bioenerg.* 36 (1995) 109.
2. K. V. Steffensen, P. Raskmark and G. F. Pedersen: FDTD calculations of the EM-field distribution in a microtiter suspension well. *Proceedings of the 244 COST meeting on "Biological effects relevant to amplitude modulated RF fields"*. Kuopio, 1995.
3. S. Kwee and P. Raskmark: The minimizing effect of electromagnetic noise on the changes in cell proliferation caused by ELF electromagnetic fields. *Proceedings of the 3rd International Congress of the European Bioelectromagnetics Association*. Nancy 1996.

MS-20-5

USE OF NON-MAMMALIAN MODEL SYSTEMS TO INVESTIGATE MECHANISMS MEDIATING BIOLOGICAL INTERACTIONS WITH ELECTROMAGNETIC FIELDS. D.M. Binnerger. Department of Biological Sciences, Florida Atlantic University, Boca Raton, Florida 33431, USA.

The question of whether power frequency (60 Hz) electromagnetic fields (EMF) present a health risk to humans remains highly controversial. While a variety of biological responses to EMF under laboratory conditions have been reported, the molecular mechanisms that mediate these phenomena remain elusive. Insight into the biochemical and genetic mechanisms that transduce EMF into biologically responsive signals is critical for clarifying the putative role of EMF in the development of certain human cancers.

The yeast *Saccharomyces cerevisiae* is an ideal model organism for studying molecular mechanisms of eukaryotic gene expression. This single-celled fungus is highly amenable to traditional genetic techniques. Additionally, recombinant DNA techniques are available for yeast which are not yet technically feasible with more complex cells, especially those of humans. While completion of the Human Genome Project remains at least 5-10 years away, the DNA sequence of the entire yeast genome is now available. Thus, experimental approaches can be designed using yeast which are not available for the other model organisms typically used in EMF studies.

Has studying yeast actually contributed to our understanding of cancer in humans? An unexpected outcome of cloning human genes has been the discovery that many cancer-related genes are very similar in all eukaryotes—from yeast to humans. Numerous human genes involved in regulating cell growth and are defective in human cancers were first identified as CDC (cell division control) genes in yeast. First identified in genetic studies using yeast, molecular clones of the yeast genes were then used to identify the homologous genes in humans. There is an ever-increasing list of human cancer genes that have structural and functional homologies in yeast.

Reported by Dr. Henry
who attended the workshop
* of Univ. of WASH., Seattle, WA